


NO-A176 545


CONFINEMENT AND REDISTRIBUTION OF CHARGES AND CURRENTS
ON A SURFACE BY EX. (U) STATE UNIV OF NEW YORK AT
BUFFALO DEPT OF CHEMISTRY H F ARNOLDUS ET AL JAN 87
UNCLASSIFIED UBUFFALO/DC/87/TR-29 N00014-86-K-0043

1/1

F/B 20/3

ML





AD-A176 545

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

R & T Code 413f0001---01

TECHNICAL REPORT No. 29

Confinement and Redistribution of Charges and Currents on a Surface
by External Fields

by

Henk F. Arnoldus, Daniel Jelski and Thomas F. George

Prepared for Publication

in

Journal of Mathematical Physics

Departments of Chemistry and Physics
State University of New York at Buffalo
Buffalo, New York 14260

January 1987

Reproduction in whole or in part is permitted for any purpose of
the United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

DTIC FILE COPY

12

DTIC
ELECTE
FEB 09 1987
S D

87 2 6 078

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/87/TR-29		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260		7b. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-86-K-0043	
8c. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NOS.	
		PROGRAM ELEMENT NO.	PROJECT NO.
		TASK NO.	WORK UNIT NO.
11. TITLE Confinement and Redistribution of Charges and Currents on a Surface by External Fields			
12. PERSONAL AUTHOR(S) Henk F. Arnoldus, Daniel Jelski and Thomas F. George			
13a. TYPE OF REPORT	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Yr., Mo., Day)	15. PAGE COUNT
16. SUPPLEMENTARY NOTATION Prepared for Publication in Journal of Mathematical Physics			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB. GR.	
		CHARGES AND CURRENTS	
		CONFINEMENT AND REDISTRIBUTION	
		SURFACE	
		EXTERNAL FIELDS	
		MAXWELL'S EQUATIONS	
		FARADAY INDUCTION	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The old problem of light scattering from a perfectly conducting surface is addressed. An electromagnetic field is incident upon the boundary, where it induces a charge and current distribution. These charges and currents emit the reflected fields. A set of equations for the charges and currents on the surface is derived by eliminating the \vec{E} - and \vec{B} -fields from Maxwell's equations with the aid of the appropriate boundary conditions. An explicit and general solution is achieved, which reveals the confinement and redistribution of the charge and the current on the surface by the external field. Expressions are obtained for the surface resolvents, or the redistribution matrices, which represent the surface geometry. Action of a surface resolvent on the incident field evaluated at the surface then yields the charge and current distributions. The Faraday induction appears as an additional contribution to the charge density. Subsequently, the reflected fields are expanded in spherical waves, which have the surface-multipole moments, and it is pointed out that charge conservation on the surface sets constraints on these moments. The results apply to arbitrarily-shaped surfaces and (con't)			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input checked="" type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson		22b. TELEPHONE NUMBER (Include Area Code) (202) 696-4410	22c. OFFICE SYMBOL

19. (con't) to any incident field. For a specific choice of the surface structure and the external field, the solutions for the charge, the current and the reflected fields are amenable to numerical evaluation.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



Journal of Mathematical Physics, in press

CONFINEMENT AND REDISTRIBUTION OF CHARGES AND CURRENTS
ON A SURFACE BY EXTERNAL FIELDS

Henk F. Arnoldus, Daniel Jelski and Thomas F. George

Departments of Chemistry and Physics
239 Fronczak Hall
State University of New York at Buffalo
Buffalo, New York 14260

PACS: 03.50, 78.20, 78.65

ABSTRACT

The old problem of light scattering from a perfectly conducting surface is addressed. An electromagnetic field is incident upon the boundary, where it induces a charge and current distribution. These charges and currents emit the reflected fields. A set of equations for the charges and currents on the surface is derived by eliminating the \underline{E} - and \underline{B} -fields from Maxwell's equations with the aid of the appropriate boundary conditions. An explicit and general solution is achieved, which reveals the confinement and redistribution of the charge and the current on the surface by the external field. Expressions are obtained for the surface resolvents, or the redistribution matrices, which represent the surface geometry. Action of a surface resolvent on the incident field evaluated at the surface, then yields the charge and current distributions. The Faraday induction appears as an additional contribution to the charge density. Subsequently, the reflected fields are expanded in spherical waves, which have the surface-multipole moments as a source. Explicit expressions are presented for the surface-multipole moments, and it is pointed out that charge conservation on the surface sets constraints on these moments. The results apply to arbitrarily-shaped surfaces and to any incident field. For a specific choice of the surface structure and the external field, the solutions for the charge, the current and the reflected fields are amenable to numerical evaluation.

I. INTRODUCTION

The study of chemistry and physics near a surface has developed rapidly during the last decade. Investigations range from classical processes like periodic deposition,¹ image formation²⁻⁴ and dispersion of plasmon waves⁵⁻²² to quantum mechanical issues as Raman scattering of intense laser light,^{23,24} atomic fluorescence near a rough surface,^{25,26} the coupling of an atomic dipole to surface polaritons²⁷ and cooperative emission processes near a conductor.²⁸ It appears, however, that besides these well-established theories, even the simplest problem --light scattering from an arbitrarily shaped surface-- is not yet completely tractable. Early approximations like the Rayleigh-Fano expansion (neglection of multiple reflections) or the small-roughness limit provide sufficient understanding of the induced effects on a boundary by incident fields, but exact solutions in the form of general expressions for the scattered fields and the surface waves are not available at present. Contemporary closed-form solutions pertain only to polarized plane waves, incident upon gratings with well-defined geometries, like square or sinusoidal wells. The results always rely on the periodicity of the surface roughness, which implies the applicability of Fourier-series expansions, or a numerical solution of the extinction theorems, as they exist in many phrasings.^{7,8,11} In this paper we consider a metallic surface, which is illuminated by an externally-applied electromagnetic field with an arbitrary time-dependence and spatial distribution. The surface is not assumed to be periodic, and our results apply equally well to a closed surface or to assemblies of surfaces, as for example a sphere near a grating. We achieve closed-form solutions of Maxwell's equations for the charge and current distributions on the surface and for the reflected fields, although at the expense of the assumption that

the metal has a perfect conductivity. The structure of the expressions is, however, quite transparent, and it might be expected that a proper account of the finite conductivity yields similar results.

II. THE FIELD EQUATIONS

The time-development and the spatial distribution of the charge density $\rho(\underline{r}, t)$, the current density $\underline{j}(\underline{r}, t)$, the electric field $\underline{E}(\underline{r}, t)$ and the magnetic field $\underline{B}(\underline{r}, t)$ are governed by Maxwell's equations. If we adopt a Fourier transform of the real fields

$$\underline{E}(\underline{r}, t) = \frac{1}{\pi} \text{Re} \int_0^{\infty} d\omega \underline{E}(\underline{r}, \omega) e^{-i\omega t}, \quad (2.1)$$

and similarly for the other three fields, then the field equations read

$$\nabla \cdot [\epsilon(\underline{r}) \underline{E}(\underline{r})] = \rho(\underline{r}), \quad (2.2)$$

$$\nabla \cdot \underline{B}(\underline{r}) = 0, \quad (2.3)$$

$$\nabla \times \underline{E}(\underline{r}) - i\omega \underline{B}(\underline{r}) = 0, \quad (2.4)$$

$$\nabla \times [\mu(\underline{r})^{-1} \underline{B}(\underline{r})] + i\omega \epsilon(\underline{r}) \underline{E}(\underline{r}) = \underline{j}(\underline{r}), \quad (2.5)$$

where we have simplified the notation by writing $\underline{E}(\underline{r})$ rather than $\underline{E}(\underline{r}, \omega)$. The frequency dependence of the fields and of $\epsilon(\underline{r})$ and $\mu(\underline{r})$ will be suppressed throughout this paper.

We shall suppose that the entire space is occupied by two kinds of media, perfect conductors and perfect insulators, which are separated by boundaries. The set of all boundaries will then be referred to as the surface. Within each medium the dielectric constant $\epsilon(\underline{r})$ and the permeability $\mu(\underline{r})$ will be assumed to be \underline{r} -independent, but across the surface $\epsilon(\underline{r})$ and $\mu(\underline{r})$ are discontinuous. Conductors are specified by a

cross product,
throughout the paper

relation like $\underline{j}(\underline{r}) = \gamma \underline{E}(\underline{r})$, $\gamma > 0$, and the assumption of perfect conductivity implies the limit $\gamma \rightarrow \infty$. Since the current density $\underline{j}(\underline{r})$ should remain finite, we obtain $\underline{E}(\underline{r}) = \underline{0}$ everywhere in the conductor. From Eq. (2.2) we then find $\rho(\underline{r}) = 0$, and Eq. (2.4) yields $\underline{B}(\underline{r}) = \underline{0}$, under the restriction $\omega \neq 0$. In this paper we will exclude the trivial static case $\omega = 0$. Finally, Eq. (2.5) gives $\underline{j}(\underline{r}) = \underline{0}$, and hence Maxwell's equations in the conductor reduce to

$$\underline{E}(\underline{r}) = \underline{0}, \underline{B}(\underline{r}) = \underline{0}, \rho(\underline{r}) = 0, \underline{j}(\underline{r}) = \underline{0}. \quad (2.6)$$

Around a point \underline{r} on the surface the fields are discontinuous. Application of Gauss' theorem on (2.2) and (2.3) and of Stokes' theorem on (2.4) and (2.5) enables us to rewrite the equations in the vicinity of the surface as

$$\underline{E}(\underline{r}^+) = \epsilon^{-1} \sigma(\underline{r}) \underline{n}(\underline{r}), \quad (2.7)$$

$$\underline{B}(\underline{r}^+) = \mu \underline{i}(\underline{r}) \times \underline{n}(\underline{r}). \quad (2.8)$$

Here $\sigma(\underline{r})$ and $\underline{i}(\underline{r})$ are the surface charge and current density, respectively, and $\underline{n}(\underline{r})$ represents the unit normal vector in \underline{r} on the surface, with the convention that it points from the conductor to the dielectric. We have introduced the notation \underline{r}^+ to indicate a point in the dielectric and close to \underline{r} . Explicitly, we write

$$\underline{r}^+ = \underline{r} + \underline{n}(\underline{r})\delta \quad \text{with } \delta \rightarrow 0. \quad (2.9)$$

We note that Eqs. (2.7) and (2.8) combine the four Maxwell equations in \underline{r} on the surface, and that they contain four unknown fields.

The dielectric is presumed to exhibit no conductivity at all, so it can be specified by $\underline{j} = \gamma \underline{E}$ with $\gamma \rightarrow 0$. This implies $\underline{j} = \underline{0}$, and from charge conservation ($\nabla \cdot \underline{j} = i\omega\rho$) we find $\rho = 0$, since we required $\omega \neq 0$. Hence, all

charges and currents, if any, are situated on the surface as $\sigma(\underline{r})$ and $\underline{i}(\underline{r})$. The electric and magnetic fields in the dielectric are generated by $\sigma(\underline{r})$ and $\underline{i}(\underline{r})$, and they contain the incident fields. This notion allows us to write Maxwell's equations for a point \underline{r} in the dielectric as

$$\rho(\underline{r}) = 0, \quad (2.10)$$

$$\underline{j}(\underline{r}) = 0, \quad (2.11)$$

$$\underline{E}(\underline{r}) - \underline{E}(\underline{r})^{\text{inc}} = \frac{-1}{4\pi\epsilon} \int dA' \sigma(\underline{r}') \nabla G(\underline{r}, \underline{r}') + \frac{i\omega\mu}{4\pi} \int dA' \underline{i}(\underline{r}') G(\underline{r}, \underline{r}'), \quad (2.12)$$

$$\underline{B}(\underline{r}) - \underline{B}(\underline{r})^{\text{inc}} = \frac{-\mu}{4\pi} \int dA' \underline{i}(\underline{r}') \times \nabla G(\underline{r}, \underline{r}'), \quad (2.13)$$

where the integrals run over the complete surface. This representation involves the Green's function of the wave equation

$$G(\underline{r}, \underline{r}') = |\underline{r} - \underline{r}'|^{-1} \exp(ik|\underline{r} - \underline{r}'|), \quad (2.14)$$

and its gradient

$$\nabla G(\underline{r}, \underline{r}') = (\underline{r} - \underline{r}') |\underline{r} - \underline{r}'|^{-3} (ik|\underline{r} - \underline{r}'| - 1) \exp(ik|\underline{r} - \underline{r}'|), \quad (2.15)$$

which contain the wave number $k = (\epsilon\mu)^{1/2}\omega$. We have to solve the set (2.12) and (2.13) for $\sigma(\underline{r})$, $\underline{i}(\underline{r})$, $\underline{E}(\underline{r})$ and $\underline{B}(\underline{r})$, and Maxwell's equations (2.7) and (2.8) on the surface can be considered as the boundary conditions.

III. ELIMINATION OF THE FIELDS

Maxwell's equations in the dielectric medium are basically two equations with four unknown fields, but we can eliminate the radiation fields $\underline{E}(\underline{r})$ and $\underline{B}(\underline{r})$ with the boundary conditions (2.7) and (2.8). To this end we take \underline{r} in (2.12) and (2.13) as \underline{r}^+ from (2.9), and then substitute the boundary values for $\underline{E}(\underline{r}^+)$ and $\underline{B}(\underline{r}^+)$. This procedure leaves us with a set of two equations for $\sigma(\underline{r})$ and $\underline{i}(\underline{r})$. The appearance of $G(\underline{r}^+, \underline{r}')$ and $\nabla_{\underline{r}^+} G(\underline{r}^+, \underline{r}')$

in the integrands of (2.12) and (2.13) is not convenient since it involves points \underline{r}^+ , which are not situated on the surface. It will turn out to be more practical to have equations in which the Green's function connects only points of the surface, rather than a point on the surface to a point in the dielectric. However, care should be exercised in replacing \underline{r}^+ by \underline{r} , because the integrals are discontinuous across the surface. If we take the limit $\underline{r}^+ \rightarrow \underline{r}$ properly (see Fig. 1), we obtain, for instance,

$$\int dA' \underline{i}(\underline{r}') \times \nabla G(\underline{r}^+, \underline{r}') = 2\pi \underline{i}(\underline{r}) \times \underline{n}(\underline{r}) + \int dA' \underline{i}(\underline{r}') \times \nabla G(\underline{r}, \underline{r}'), \quad (3.1)$$

and we observe that replacing \underline{r}^+ by \underline{r} requires that we should add the term $2\pi \underline{i}(\underline{r}) \times \underline{n}(\underline{r})$. It was already pointed out by Maradudin²⁹ that integrals of this kind appear to have a finite contribution from a single point. This feature can however also be regarded as resulting from the discontinuity of the fields across the surface. Critical comments on this issue have also been made by Agarwal¹² in a slightly different context.

With the explicit forms of $G(\underline{r}, \underline{r}')$ and its gradient, as they are given by (2.14) and (2.15), we can evaluate the limit $\underline{r}^+ \rightarrow \underline{r}$, which yields the set of equations

$$\begin{aligned} \sigma(\underline{r}) \underline{n}(\underline{r}) = & \frac{-1}{2\pi} \int dA' \sigma(\underline{r}') \nabla G(\underline{r}, \underline{r}') + \frac{i\omega\epsilon\mu}{2\pi} \int dA' \underline{i}(\underline{r}') G(\underline{r}, \underline{r}') \\ & + 2E(\underline{r})^{\text{inc}}, \end{aligned} \quad (3.2)$$

$$\underline{i}(\underline{r}) \times \underline{n}(\underline{r}) = \frac{-1}{2\pi} \int dA' \underline{i}(\underline{r}') \times \nabla G(\underline{r}, \underline{r}') + 2\mu^{-1} \underline{B}(\underline{r})^{\text{inc}}, \quad (3.3)$$

for $\sigma(\underline{r})$ and $\underline{i}(\underline{r})$. We can write $\sigma(\underline{r})$ and $\underline{i}(\underline{r})$ in the integrands as

$$\sigma(\underline{r}) = \underline{n}(\underline{r}) \cdot (\sigma(\underline{r}) \underline{n}(\underline{r})), \quad (3.4)$$

$$\underline{i}(\underline{r}) = \underline{n}(\underline{r}) \times (\underline{i}(\underline{r}) \times \underline{n}(\underline{r})), \quad (3.5)$$

since $\underline{i}(\underline{r})$ is parallel to the surface, which shows that Eqs. (3.2) and (3.3) are essentially a set of equations for the vector fields $\sigma(\underline{r})\underline{n}(\underline{r})$ and $\underline{i}(\underline{r})\underline{x}\underline{n}(\underline{r})$ on the surface.

Equation (3.3) for $\underline{i}(\underline{r})\underline{x}\underline{n}(\underline{r})$ has the form of an inhomogeneous Fredholm equation of the second kind, where the external field $2\mu^{-1}\underline{B}(\underline{r})^{\text{inc}}$ is the inhomogeneity. In the same fashion, Eq. (3.2) has $2\varepsilon\underline{E}(\underline{r})^{\text{inc}}$ (and the current term) as an inhomogeneous part. Hence the incident fields can be regarded as the source terms of these equations. In this sense $\sigma(\underline{r}) \neq 0$ and $\underline{i}(\underline{r}) \neq \underline{0}$ are a result of the presence of the driving field, so the charges and the currents are confined on the surface by the field. If there is a net charge on the surface, this mechanism might also be conceived as a redistribution process. Equations (3.2) and (3.3) resemble the extinction theorem for the analogous problem of scattering of an incident field from a dielectric grating. The extinction theorem is, however, a homogeneous equation, and its solvability condition is equivalent to the dispersion relation for surface polaritons.

IV. REPRESENTATION OF THE SURFACE

Ordinary Fredholm equations are single-variable equations for a function on the complex plane, and they can be solved by an expansion of the function onto a suitable complete set. Our equations for $\sigma(\underline{r})\underline{n}(\underline{r})$ and $\underline{i}(\underline{r})\underline{x}\underline{n}(\underline{r})$ are three-dimensional and surface-related equations for a vector field, so we have to modify the standard techniques slightly. In order to accomplish this, we introduce spherical coordinates (r, θ, ϕ) with respect to an arbitrary origin, and we will abbreviate the direction θ, ϕ by the single variable Ω . Then the assembly of all points \underline{r} , which constitute the surface, can be represented by a set of functions $\xi(\Omega)_\lambda$. The $\xi(\Omega)_\lambda$ will indicate the distance from the origin to a point \underline{r} on the surface, in the

direction Ω , while the subscript λ accounts for the multiplicity (see Fig. 2). In this fashion, the surface is divided in regions, numbered by λ , where its shape is defined by a function $\xi(\Omega)_\lambda$, which determines uniquely the spherical coordinates $(r, \theta, \phi) = (\xi(\theta, \phi)_\lambda, \theta, \phi)$ of a point \underline{r} in this region. The shape-functions $\xi(\Omega)_\lambda$ will be assumed to be given, and therefore we can represent a point on the surface by its surface-coordinates (λ, Ω) rather than by its spherical coordinates (r, Ω) . We will use λ as a subscript and Ω as a variable.

The measure $dA(\Omega)_\lambda$ and the direction $\underline{n}(\Omega)_\lambda$ of the surface at a given point (λ, Ω) are fixed by its shape $\xi(\Omega)_\lambda$. For instance, the infinitesimal surface area at (λ, Ω) is given by

$$dA(\Omega)_\lambda = f(\Omega)_\lambda d\Omega \quad (4.1)$$

with

$$f(\Omega)_\lambda = \xi(\Omega)_\lambda \left\{ \xi(\Omega)_\lambda^2 + \left(\frac{\partial}{\partial \theta} \xi(\Omega)_\lambda \right)^2 + \frac{1}{\sin^2 \theta} \left(\frac{\partial}{\partial \phi} \xi(\Omega)_\lambda \right)^2 \right\}^{1/2}, \quad (4.2)$$

in terms of the infinitesimal surface area $d\Omega = \sin \theta d\theta d\phi$ of the unit sphere. Hence the function $f(\Omega)_\lambda$ accounts for the deviation of the surface curvature from the curvature of a sphere, and with the aid of (4.1) we can transform a surface-integral over the region λ into an integration over a part of the unit sphere. We note that not every direction Ω for a given λ corresponds to a point on the surface. It will turn out to be convenient to extend the definition (4.1) of $f(\Omega)_\lambda$ as

$$f(\Omega)_\lambda = 0 \quad \text{if } \Omega \text{ does not correspond to a point on} \\ \text{the surface in region } \lambda. \quad (4.2)$$

Then we can write the surface integrals as

$$\int d\Lambda \dots = \sum_{\lambda} \int d\Omega f(\Omega)_{\lambda} \dots, \quad (4.3)$$

where the integrals now run over the complete unit sphere for every λ . This construction will enable us to apply the general theory of expanding vector fields on a sphere.

V. EXPANSION OF THE FIELDS

Since we are using spherical coordinates, the spherical harmonics $Y(\Omega)_{\ell m}$ supply a suitable complete set on the unit sphere for an expansion of the magnitude of a vector field. The direction of a vector will be expanded onto a space-fixed set of three unit vectors, denoted by $\underline{e}_{-\tau}$, which is, for instance, the Cartesian set $\underline{e}_x, \underline{e}_y, \underline{e}_z$ or the spherical set $\underline{e}_{+1}, \underline{e}_0, \underline{e}_{-1}$. Then the vector fields $Y(\Omega)_{\ell m} \underline{e}_{-\tau}$ constitute a complete set on the unit sphere for an expansion of an arbitrary vector field.

It is our aim to solve Eqs. (3.2) and (3.3) for $\sigma(\underline{r})\underline{n}(\underline{r})$ and $\underline{i}(\underline{r})\underline{x}\underline{n}(\underline{r})$. We thus start with an expansion of these fields,

$$f(\Omega)_{\lambda} \sigma(\Omega)_{\lambda} \underline{n}(\Omega)_{\lambda} = \sum_{\ell m \tau} S_{\ell m \tau \lambda} Y_{\ell m}(\Omega) \underline{e}_{-\tau}, \quad (5.1)$$

$$f(\Omega)_{\lambda} \underline{i}(\Omega)_{\lambda} \underline{x} \underline{n}(\Omega)_{\lambda} = \sum_{\ell m \tau} I_{\ell m \tau \lambda} Y_{\ell m}(\Omega) \underline{e}_{-\tau}, \quad (5.2)$$

and note that we have included a factor $f(\Omega)_{\lambda}$ on the left-hand side. This is necessary, since otherwise the left-hand side of Eqs. (5.1) and (5.2) would not be properly defined for every Ω . The driving, incident fields $\underline{E}(\underline{r})^{inc}, \underline{B}(\underline{r})^{inc}$ in Eqs. (3.2) and (3.3) enter only through their value on the surface, so that we can expand them on the surface set according to

$$f(\Omega)_{\lambda} \underline{E}(\Omega)_{\lambda}^{inc} = \sum_{\ell m \tau} E_{\ell m \tau \lambda} Y_{\ell m}(\Omega) \underline{e}_{-\tau}, \quad (5.3)$$

$$f(\Omega)_{\lambda} \underline{B}(\Omega)_{\lambda}^{inc} = \sum_{\ell m \tau} B_{\ell m \tau \lambda} Y_{\ell m}(\Omega) \underline{e}_{-\tau}. \quad (5.4)$$

The expansion coefficients for the incident fields then follow from the inverse relation

$$E_{\ell m \tau \lambda} = \int d\Omega f(\Omega)_{\lambda} E(\Omega)_{\lambda}^{\text{inc}} \cdot e_{-\tau}^* Y(\Omega)_{\ell m}^* \quad (5.5)$$

$$B_{\ell m \tau \lambda} = \int d\Omega f(\Omega)_{\lambda} B(\Omega)_{\lambda}^{\text{inc}} \cdot e_{-\tau}^* Y(\Omega)_{\ell m}^* \quad (5.6)$$

and the appearance of $f(\Omega)_{\lambda}$ in the integrands reflects that we actually have integrals over region λ of the surface. This illustrates that $f(\Omega)_{\lambda} Y(\Omega)_{\ell m}^* e_{-\tau}^*$ can be considered as a complete surface set for the expansion of a vector field on the surface. Note that we allow $e_{-\tau}$ to be complex, which is the case for a spherical set.

VI. THE CHARGE AND THE CURRENT DISTRIBUTIONS

It is straightforward to rewrite Eqs. (3.2) and (3.3) for $\sigma(\underline{r})\underline{n}(\underline{r})$ and $\underline{i}(\underline{r})\underline{x}\underline{n}(\underline{r})$ in terms of their expansion coefficients. We obtain

$$\sum_{\ell' m' \tau' \lambda'} (R_{\ell m \tau \lambda, \ell' m' \tau' \lambda'}^{(2)} - \delta_{\ell \ell'} \delta_{\tau \tau'} \delta_{\lambda \lambda'}) S_{\ell' m' \tau' \lambda'} = i\omega \epsilon \mu \sum_{\ell' m' \tau' \lambda'} R_{\ell m \tau \lambda, \ell' m' \tau' \lambda'}^{(3)} I_{\ell' m' \tau' \lambda'} - 2\epsilon E_{\ell m \tau \lambda} \quad (6.1)$$

$$\sum_{\ell' m' \tau' \lambda'} (R_{\ell m \tau \lambda, \ell' m' \tau' \lambda'}^{(1)} - \delta_{\ell \ell'} \delta_{\tau \tau'} \delta_{\lambda \lambda'}) I_{\ell' m' \tau' \lambda'} = -2\mu^{-1} B_{\ell m \tau \lambda} \quad (6.2)$$

which are two coupled inhomogeneous linear equations for the surface charge $S_{\ell m \tau \lambda}$ and the surface current $I_{\ell m \tau \lambda}$. The expansion coefficients $E_{\ell m \tau \lambda}$ and $B_{\ell m \tau \lambda}$ for the external fields are supposed to be given. The set (6.1) and (6.2) also involves three R-matrices, with matrix elements

$$R_{\ell m \tau \lambda, \ell' m' \tau' \lambda'}^{(1)} = \frac{-1}{2\pi} \int d\Omega \int d\Omega' f(\Omega)_{\lambda} Y(\Omega)_{\ell m}^* Y(\Omega')_{\ell' m'} \cdot \{e_{-\tau}^* x(n(\Omega')_{\lambda}) x_{e_{-\tau}}\} \cdot \nabla G(\Omega, \Omega')_{\lambda \lambda'}, \quad (6.3)$$

$$R_{\ell m \tau \lambda, \ell' m' \tau' \lambda'}^{(2)} = \frac{-1}{2\pi} \int d\Omega \int d\Omega' f(\Omega) Y_{\ell m}(\Omega) Y_{\ell' m'}^*(\Omega') \cdot (\underline{n}(\Omega')_{\lambda'} \cdot \underline{e}_{\tau'}^*) \underline{e}_{\tau}^* \cdot \nabla G(\Omega, \Omega')_{\lambda \lambda'}, \quad (6.4)$$

$$R_{\ell m \tau \lambda, \ell' m' \tau' \lambda'}^{(3)} = \frac{-1}{2\pi} \int d\Omega \int d\Omega' f(\Omega) Y_{\ell m}(\Omega) Y_{\ell' m'}^*(\Omega') \cdot \underline{e}_{\tau}^* (\underline{n}(\Omega')_{\lambda'} \cdot \underline{x} \underline{e}_{\tau'}) G(\Omega, \Omega')_{\lambda \lambda'}, \quad (6.5)$$

where we have written $G(\Omega, \Omega')_{\lambda \lambda'}$ for the Green's function, which connects the points (λ, Ω) and (λ', Ω') of the surface. We emphasize that these R-matrices depend only on the geometry of the surface, and not on the external fields. Prescription of the shape of the surface determines the R-matrices. Recall, however, that the R-matrices depend on the frequency ω through the Green's function, but this is merely a parametric dependence and independent of the external field.

The expansion coefficients $S_{\ell m \tau \lambda}$ can always be arranged in a one-dimensional array, considered as a vector, and similarly $R^{(1)}$, $R^{(2)}$ and $R^{(3)}$ can be regarded as two-dimensional matrices. Then we can write (6.1) and (6.2) as

$$(R^{(2)} - 1)S = i\omega \epsilon \mu R^{(3)} I - 2\epsilon E, \quad (6.6)$$

$$(R^{(1)} - 1)I = -2\mu^{-1}B, \quad (6.7)$$

where we have also adopted a vector representation for the driving fields. The solution of (6.6) and (6.7) is immediately found to be

$$S = \frac{2\epsilon}{1 - R^{(2)}} \left\{ E - i\omega R^{(3)} \frac{1}{1 - R^{(1)}} B \right\}, \quad (6.8)$$

$$I = \frac{2\mu^{-1}}{1 - R^{(1)}} B, \quad (6.9)$$

which expresses the charge density S and the current density I explicitly in the externally applied fields E and B and the surface-shape matrices $R^{(1)}$, $R^{(2)}$ and $R^{(3)}$.

For vanishing external fields, e.g., $E = 0$ and $B = 0$, the charge and current distributions also vanish, as can be seen explicitly from Eqs. (6.8) and (6.9). Hence the charges and currents are indeed confined to the surface by the external fields. Remember that we have excluded the static case $\omega = 0$, for which we can have charges on a surface without external fields. Furthermore, we can identify the resolvents $(1-R^{(2)})^{-1}$ and $(1-R^{(1)})^{-1}$ as the operators which account for the redistribution of the charges and currents, respectively, as resulting from the Lorentz force between charges and between currents. The coupling of charges and currents, which is the Faraday induction, is incorporated in the $R^{(3)}$ -matrix.

VII. THE REFLECTED FIELDS

The incident field induces charges and currents on the surface, and these oscillating charges and currents emit radiation, which are the reflected fields. In this section we express these fields in terms of the expansion coefficients $S_{\ell m \tau \lambda}$ and $I_{\ell m \tau \lambda}$, as they are given explicitly in the previous section.

In Eq. (2.12) we expressed the reflected electric field $\underline{E}(\underline{r}) - \underline{E}(\underline{r})^{\text{inc}}$ in terms of $\sigma(\underline{r})$ and $\underline{i}(\underline{r})$, and similarly $\underline{B}(\underline{r}) - \underline{B}(\underline{r})^{\text{inc}}$ in terms of $\underline{i}(\underline{r})$. With (3.4) and (3.5) we can rewrite these equations in a way that $\sigma(\underline{r})\underline{n}(\underline{r})$ and $\underline{i}(\underline{r})\underline{x}\underline{n}(\underline{r})$ are the source fields, and then we can apply (5.1) and (5.2) in order to find an expansion on the spherical set. However, the resulting expressions are not transparent, since they will involve the Green's function and its gradient. In order to achieve a more comprehensible result, we expand the Green's function on the spherical set. We write²³

$$G(\underline{r}, \underline{r}') = 4\pi i k \sum_{\ell m} h_{\ell}^{(1)}(k\xi(\Omega'))_{\lambda} Y(\Omega')_{\ell m} j_{\ell}(kr)_{\ell} Y(\Omega)_{\ell m}^*, \quad (7.1)$$

where $h_{\ell}^{(1)}$ and j_{ℓ} are spherical Bessel functions. Here the convention is that we choose the origin of our coordinate system in the dielectric, and in such a way that the inequality

$$\xi(\Omega)_{\lambda} > r \quad (7.2)$$

holds for every (λ, Ω) . The vector \underline{r} is the position in the dielectric, where we wish to evaluate the reflected fields. The expansion coefficients $S_{\ell m \tau \lambda}$ and $I_{\ell m \tau \lambda}$ depend on the position of the origin, so both the charge and current distributions and the reflected fields must be evaluated with respect to the same coordinate system. Furthermore, restriction (7.2) must hold in order to apply the series expansion (7.1) of the Green's function. For a given \underline{r} , this can always be arranged.

The solution for the fields can be cast in an appealing form by the introduction of the source-term vectors

$$\underline{S}_{\ell m}^{(\lambda)} = \sum_{\tau} S_{\ell m \tau \lambda} \underline{e}_{\tau} \quad (7.3)$$

$$\underline{I}_{\ell m}^{(\lambda)} = \sum_{\tau} I_{\ell m \tau \lambda} \underline{e}_{\tau}. \quad (7.4)$$

In view of (5.1) and (5.2), these $\underline{S}_{\ell m}^{(\lambda)}$ and $\underline{I}_{\ell m}^{(\lambda)}$ are just the expansion coefficients of $f(\Omega)_{\lambda} \sigma(\Omega)_{\lambda} \underline{n}(\Omega)_{\lambda}$ and $f(\Omega)_{\lambda} i(\Omega)_{\lambda} \underline{x} \underline{n}(\Omega)_{\lambda}$ after an expansion of these fields onto the set of spherical harmonics, but without a decomposition along the basis vectors \underline{e}_{τ} . Furthermore, we define the vector

$$\underline{P}_{\ell m, \ell' m'}^{(\lambda)} = -i \int d\Omega f(\Omega)_{\lambda} Y(\Omega)_{\ell m} Y(\Omega)_{\ell' m'} h_{\ell}^{(1)}(k\xi(\Omega))_{\lambda} \underline{n}(\Omega)_{\lambda}, \quad (7.5)$$

which is a surface integral over the region λ . It is the integrated normal vector $\underline{n}(\Omega)_{\lambda}$ times the appropriate weight-functions. This vector $\underline{P}_{\ell m, \ell' m'}^{(\lambda)}$ depends only on the shape of the surface. After these preliminary

definitions, we can write for the reflected fields

$$\begin{aligned} \underline{E}(\underline{r}) - \underline{E}(\underline{r})^{inc} = & \frac{k}{\epsilon} \sum_{\ell m \ell' m' \lambda} \underline{p}_{\ell m, \ell' m'}^{(\lambda)} \cdot \underline{S}_{\ell m}^{(\lambda)} \nabla j(kr)_{\ell} Y(\Omega)_{\ell' m'}^* \\ & - i\omega\mu k \sum_{\ell m \ell' m' \lambda} \underline{p}_{\ell m, \ell' m'}^{(\lambda)} \times \underline{I}_{\ell m}^{(\lambda)} j(kr)_{\ell} Y(\Omega)_{\ell' m'}^*, \end{aligned} \quad (7.6)$$

$$\underline{B}(\underline{r}) - \underline{B}(\underline{r})^{inc} = \mu k \sum_{\ell m \ell' m' \lambda} (\underline{p}_{\ell m, \ell' m'}^{(\lambda)} \times \underline{I}_{\ell m}^{(\lambda)}) \nabla j(kr)_{\ell} Y(\Omega)_{\ell' m'}^*. \quad (7.7)$$

These explicit expressions for the fields that are emitted by the surface charge and current distributions exhibit a clear separation between the source terms $\underline{S}_{\ell m}^{(\lambda)}$ and $\underline{I}_{\ell m}^{(\lambda)}$ and the redistribution, due to the surface geometry, which is accounted for by the vector $\underline{p}_{\ell m, \ell' m'}^{(\lambda)}$. The spatial distribution is represented as an expansion in the spherical waves $j(kr)_{\ell} Y(\Omega)_{\ell m}^*$ and $\nabla j(kr)_{\ell} Y_{\ell m}(\Omega)^*$.

VIII. SURFACE MULTIPOLES

We can elucidate the significance of the expansions (7.6) and (7.7) for the reflected fields by the introduction of the surface multipoles. To this end we define the multipolar moments of the charge and the current distributions as

$$\underline{C}_{\ell m} = \frac{k}{\epsilon} \sum_{\ell' m' \lambda} \underline{p}_{\ell' m', \ell m}^{(\lambda)} \cdot \underline{S}_{\ell' m'}^{(\lambda)}, \quad (8.1)$$

$$\underline{J}_{\ell m} = \mu k \sum_{\ell' m' \lambda} \underline{p}_{\ell' m', \ell m}^{(\lambda)} \times \underline{I}_{\ell' m'}^{(\lambda)}, \quad (8.2)$$

where $\underline{C}_{\ell m}$ is a scalar and $\underline{J}_{\ell m}$ is a vector. These multipolar moments represent the charge and current distribution of the complete surface, not just in one region λ . The emitted fields now attain the form

$$\underline{E}(\underline{r}) - \underline{E}(\underline{r})^{inc} = \sum_{\ell m} \underline{C}_{\ell m} \nabla j(kr)_{\ell} Y(\Omega)_{\ell m}^* - i\omega \sum_{\ell m} \underline{J}_{\ell m} j(kr)_{\ell} Y(\Omega)_{\ell m}^*, \quad (8.3)$$

$$\underline{B}(\underline{r}) - \underline{B}(\underline{r})^{\text{inc}} = \sum_{\ell m} \underline{J}_{\ell m} \times \nabla j(kr)_{\ell} Y(\Omega)_{\ell m}^* \quad (8.4)$$

which greatly resembles the multipole expansion of the fields emitted by a charge and current distribution in a restricted region of space. The distinction is of course that the source terms $C_{\ell m}$ and $\underline{J}_{\ell m}$ here gain contributions from everywhere in space, rather than from a localized area. This results effectively in an exchange of the spherical Bessel function $h^{(1)}(kr)_{\ell}$ with $j(kr)_{\ell}$ in the expansion of the Green's function.

The surface multipolar moments $C_{\ell m}$ and $\underline{J}_{\ell m}$ are not independent. From the fact that the fields obey Maxwell's equations, as they do by construction, it follows that they are subject to some constraints. From $\nabla \cdot (\underline{E}(\underline{r}) - \underline{E}(\underline{r})^{\text{inc}}) = 0$ we readily derive the relation

$$\begin{aligned} i\sqrt{\epsilon\mu}C_{\ell m} = & \frac{\sqrt{\ell}}{\sqrt{2\ell-1}} \sum_{\mu=-(\ell-1)}^{\ell-1} \sum_{\tau} (\ell m 1 \tau | \ell-1 \mu) \underline{J}_{\ell-1, \mu} \cdot \underline{e}_{\tau}^* \\ & + \frac{\sqrt{\ell+1}}{\sqrt{2\ell+3}} \sum_{\mu=-(\ell+1)}^{\ell+1} \sum_{\tau} (\ell m 1 \tau | \ell+1 \mu) \underline{J}_{\ell+1, \mu} \cdot \underline{e}_{\tau}^* \end{aligned} \quad (8.5)$$

for a spherical basis set \underline{e}_{τ} . Here $(\ell m 1 \tau | \ell \pm 1 \mu)$ denotes a Clebsch-Gordan coefficient. The constraint (8.5) can be considered as the surface-integrated form of charge conservation ($\nabla \cdot \underline{j} = i\omega\rho$) for the surface charge density $\sigma(\underline{r})$.

IX. CONCLUSIONS

We have studied the charge and current distributions on the boundary of a perfect conductor with a dielectric, as they are confined and redistributed there by an externally applied electromagnetic field. The surface was allowed to have an arbitrary shape, and we did not impose any

periodicity condition. We obtained closed-form and exact expressions for $\sigma(\underline{r})$ and $\underline{i}(\underline{r})$ everywhere on the surface. This was accomplished by deriving a set of inhomogeneous Fredholm equations of the second kind for $\sigma(\underline{r})\underline{n}(\underline{r})$ and $\underline{i}(\underline{r})\underline{x}\underline{n}(\underline{r})$ from Maxwell's equations, and subsequently solving these equations by an expansion on a discrete spherical set of basis vector functions. The solution involves surface-structure matrices, the R -matrices, which are independent of the incident field. It appears that an operation of a resolvent $(1-R)^{-1}$ on the vector representation of the impinging field on the surface yields the charge and current distributions. The Faraday induction between the \underline{E} - and the \underline{B} -fields gives rise to a coupling between the equations for $\sigma(\underline{r})\underline{n}(\underline{r})$ and $\underline{i}(\underline{r})\underline{x}\underline{n}(\underline{r})$, and it was accounted for by the matrix $R^{(3)}$.

Next, the structure of the fields, which are emitted by the oscillating charges and currents, was examined. The solution was cast in the form of a spherical multipolar expansion, and the multipolar moments were identified explicitly in terms of the solutions for $\sigma(\underline{r})$ and $\underline{i}(\underline{r})$. The effect of the surface geometry could be incorporated entirely by the application of a surface-integrated normal-direction matrix $p_{\ell m, \ell' m'}^{(\lambda)}$. In addition, it was shown that the multipolar moments for the charge and current distributions are related, which reflects the charge conservation on the surface.

ACKNOWLEDGEMENTS

This research was supported by the Office of Naval Research and the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009. The United States Government is authorized to reproduce and distribute reprints notwithstanding any copyright notation hereon.

REFERENCES

1. S.R.J. Brueck and D.J. Ehrlich, Phys. Rev. Lett. 48, 1678 (1982).
2. A.G. Ramm and M.A. Fiddy, Opt. Commun. 56, 8 (1985).
3. P.J. Feibelman, Phys. Rev. B 12, 1319 (1975).
4. P.J. Feibelman, Phys. Rev. B 12, 4282 (1975).
5. H.J. Juranek, Z. Physik 233, 324 (1970).
6. J.P. Marton and J.R. Lemon, Phys. Rev. B 4, 271 (1971).
7. J.J. Sein, Opt. Commun. 14, 157 (1975).
8. G.S. Agarwal, Opt. Commun. 14, 161 (1975).
9. A.A. Maradudin and D.L. Mills, Phys. Rev. B 11, 1392 (1975).
10. V. Celli, A. Marvin and F. Toigo, Phys. Rev. B 11, 1779 (1975).
11. A. Marvin, F. Toigo and V. Celli, Phys. Rev. B 11, 2777 (1975).
12. G.S. Agarwal, Phys. Rev. B 14, 846 (1976).
13. F. Toigo, A. Marvin, V. Celli and N.R. Hill, Phys. Rev. B 15, 5618 (1977).
14. N. Garcia and N. Cabrera, Phys. Rev. B 18, 576 (1978).
15. B. Laks, D.L. Mills and A.A. Maradudin, Phys. Rev. B 23, 4965 (1981).
16. N.E. Glass and A.A. Maradudin, Phys. Rev. B 24, 595 (1981).
17. P. Sheng, R.S. Stepleman and P.N. Sanda, Phys. Rev. B 26, 2907 (1982).
18. N. Garcia and A.A. Maradudin, Opt. Commun. 45, 301 (1983).
19. N. Garcia, Opt. Commun. 45, 307 (1983).
20. M. Weber and D.L. Mills, Phys. Rev. B 27, 2698 (1983).
21. N.E. Glass, M. Weber and D.L. Mills, Phys. Rev. B 29, 6548 (1984).
22. K.T. Lee and T.F. George, Phys. Rev. B 31, 5106 (1985).
23. S.S. Jha, J.R. Kirtley and J.C. Tsang, Phys. Rev. B 22, 3973 (1980).
24. M. Nevière and R. Reinisch, Phys. Rev. B 26, 5403 (1982).

25. X.Y. Huang, K.T. Lee and T.F. George, J. Chem. Phys., in press.
26. D. Agassi, Phys. Rev. B 32, (1985), in press.
27. G.S. Agarwal and C.V. Kunasz, Phys. Rev. B 26, 5832 (1982).
28. K.C. Liu and T.F. George, Phys. Rev. B 32, 3622 (1985).
29. A.A. Maradudin, in Surface Polaritons, edited by V.M. Agranovich and A.A. Maradudin (North-Holland, Amsterdam, 1982), pp. 405.
30. P.M. Morse and H. Feshback, Methods of Theoretical Physics, (McGraw-Hill, New York, 1953), pp. 887.

FIGURE CAPTIONS

Figure 1. Geometry for the evaluation of the limit $\underline{r}^+ \rightarrow \underline{r}$. Around \underline{r} on the surface, we divide the surface into an infinitesimal circle of radius R , and the rest of the surface. Then the integrals are split up accordingly. The normal vector points from the surface into the dielectric and is multiplied by $\delta > 0$. We denoted $\underline{r}^+ - \underline{r}'$ by \underline{a} and $\underline{r} - \underline{r}'$ by \underline{b} . The limit $\underline{r}^+ \rightarrow \underline{r}$ implies $R \gg \delta > 0$ and $R \rightarrow 0$. It appears that an integral over the small circle remains finite whenever the gradient of the Green's function occurs in the integrand. This gives rise to the first term on the right-hand side of Eq. (3.1).

Figure 2. Illustration of the surface multiplicity. From the origin O in the direction Ω , we find points on the surface which have a distance $\xi(\Omega)_1, \xi(\Omega)_2, \dots$ to O . Therefore, a description of the surface in spherical coordinates requires a set of functions $\xi(\Omega)_\lambda$.

Fig. 1





Fig. 2



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. A. Reisman
Microelectronics Center of North Carolina
Research Triangle Park, North Carolina
27709

Dr. M. Grunze
Laboratory for Surface Science and
Technology
University of Maine
Orono, Maine 04469

Dr. J. Butler
Naval Research Laboratory
Code 6115
Washington D.C. 20375-5000

Dr. L. Interante
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Irvin Heard
Chemistry and Physics Department
Lincoln University
Lincoln University, Pennsylvania 19352

Dr. K.J. Klaubunde
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookeville, Tennessee 38501

Dr. D. DiLella
Chemistry Department
George Washington University
Washington D.C. 20052

Dr. R. Reeves
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Steven M. George
Stanford University
Department of Chemistry
Stanford, CA 94305

Dr. Mark Johnson
Yale University
Department of Chemistry
New Haven, CT 06511-8118

Dr. W. Knauer
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. Murday
Naval Research Laboratory
Code 6170
Washington, D.C. 20375-5000

Dr. J. B. Hudson
Materials Division
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne
Chemistry Department
Northwestern University
Evanston, Illinois 60637

Dr. J. M. White
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. D. E. Harrison
Department of Physics
Naval Postgraduate School
Monterey, California 93940

Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria
Electrical Engineering Department
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green
Quantum Surface Dynamics Branch
Code 3817
Naval Weapons Center
China Lake, California 93555

Dr. A. Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02912

Dr. S. L. Bernasek
Department of Chemistry
Princeton University
Princeton, New Jersey 08544

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Richard Colton
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Dan Pierce
National Bureau of Standards
Optical Physics Division
Washington, D.C. 20234

Dr. R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer
Materials Characterization Lab.
General Electric Company
Schenectady, New York 22217

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Ronald Lee
R301
Naval Surface Weapons Center
White Oak
Silver Spring, Maryland 20910

Dr. Paul Schoen
Code 6190
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. John T. Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene
Code 5230
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda
University of Pittsburgh
Chemistry Building
Pittsburg, PA 15260

Dr. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH
UNITED KINGDOM

Dr. H. Tachikawa
Chemistry Department
Jackson State University
Jackson, Mississippi 39217

Dr. John W. Wilkins
Cornell University
Laboratory of Atomic and
Solid State Physics
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis
Department of Physics
University of California
Irvine, California 92664

Dr. D. Ramaker
Chemistry Department
George Washington University
Washington, D.C. 20052

Dr. J. C. Hemminger
Chemistry Department
University of California
Irvine, California 92717

Dr. T. F. George
Chemistry Department
University of Rochester
Rochester, New York 14627

Dr. G. Rubloff
IBM
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. Horia Metiu
Chemistry Department
University of California
Santa Barbara, California 93106

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. P. Hansma
Department of Physics
University of California
Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser
Department of Chemistry
University of Richmond
Richmond, Virginia 23173

Dr. R. W. Plummer
Department of Physics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. Steckl
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. G.H. Morrison
Department of Chemistry
Cornell University
Ithaca, New York 14853

END

3-87

Dtic